

In order to have the best analyse and avoid contamination, please follow strictly these steps:

- Urine is preferably collected in the morning and/or after at least 4 hours after the previous urination.
- Before starting antibiotic treatment



**1 - Wash your hands with water and soap; unscrew the lid of the container and place it on a flat surface**



**2 - Clean your genital area with water and soap, or with Dakin using sterile compress**




**3- Pass the first urine into the toilet**



**4 - Collect the next stream into the sterile container (mid-stream urine)**



**5 - Close the container securely**




**6 - Wash your hands and return the container to the nurse**

ដើម្បីធានាថាការវិភាគមានភាពត្រឹមត្រូវ និងជៀសវាង ការចម្លងមេរោគពីរខាងក្រៅ សូមលោកអ្នកអនុវត្តតាម ជំហាន ខាងក្រោមអោយបានត្រឹមត្រូវ៖

- ការយកសំណាកត្រូវធ្វើនៅពេលព្រឹក ឬបន្ទាប់ពីការទប់នោមយ៉ាងហោចណាស់ ៤ ម៉ោង។
- មុនពេលចាប់ផ្តើមការប្រើប្រាស់ថ្នាំផ្សេងៗ



1 - លាងដៃជាមួយសាប៊ូឲ្យបានស្អាត និងបើកគម្របកំប៉ុង រួចទុកនៅកន្លែងរាបស្មើ



2 - សម្អាតតំបន់ប្រដាប់ភេទ ដោយ ប្រើ ទឹក និងសាប៊ូ ឬទឹកសម្លាប់ មេរោគ Dakin



3 - នោមទឹកនោមដំបូងចោលបន្តិច ទៅក្នុងបង្គន់



4- ប្រមូលទឹកនោមផ្នែកកណ្តាល ដាក់ក្នុងកំប៉ុង




5 - មូលគម្របកំប៉ុងឲ្យជិត



6- លាងដៃរបស់អ្នក ហើយប្រគល់កំប៉ុង ទៅឲ្យគិលានុបដ្ឋាយិកា

FIT-R1-002 - version 3 du 09/04/2026 / (02/06/2026)